

EXTENSIONS OF VERMA MODULES

R. VIRK

Fix a finite dimensional semisimple Lie algebra over \mathbf{C} , and a Borel subalgebra. It is a long standing problem to determine the extension groups between Verma modules in the category \mathcal{O} of [BGG]. No general formula, even conjectural, is known.

Let W be the Weyl group and $w_0 \in W$ the longest element. For $w \in W$ write Δ_w for the Verma module of highest weight $w^{-1}w_0 \cdot 0$ (dot action). The purpose of this note is to give a recursive formula (Corollary 5) for $\dim \text{Ext}^1(\Delta_v, \Delta_w)$ (throughout ' $\text{Ext}^\bullet = \text{Ext}_{\mathcal{O}}^\bullet$ ').

Write X for the flag variety associated to our Lie algebra. For each $w \in W$, we have a Schubert cell $C_w \subseteq X$ and an opposite Schubert cell $C^w \subseteq X$. Let $\ell: W \rightarrow \mathbf{Z}_{\geq 0}$ denote the length function and \leq the Bruhat order. If $v \leq w$, then $C^v \cap C_w$ is affine and smooth of dimension $\ell(w) - \ell(v)$ (see [R]). Further, $C^v \cap C_v = \text{pt}$; and if $v \not\leq w$, then $C^v \cap C_w = \emptyset$.

By [RSW, Proposition 4.2.1] and [BGS, Proposition 3.5.1]):

$$\text{Ext}^\bullet(\Delta_v, \Delta_w) = H_c^{\bullet + \ell(w) - \ell(v)}(C^v \cap C_w), \quad \text{for all } v, w \in W,$$

where H_c^* denotes compactly supported cohomology (with \mathbf{C} -coefficients). The cohomology $H_c^*(C^v \cap C_w)$ comes equipped with a canonical (rational) Hodge structure which is respected by the usual long exact sequences. Consequently, we can and will view the extensions $\text{Ext}^\bullet(\Delta_v, \Delta_w)$ as Hodge structures. Denote the trivial 1-dimensional Hodge structure by \mathbf{Q}^H , and write (n) for the n -th Tate twist.

The following Proposition is attributed to V. Deodhar in [RSW] (see [RSW, Lemma 4.3.1] for a proof).

Proposition 1. *Let $v, w \in W$ with $v \leq w$. Let $s \in W$ be a simple reflection such that $ws < w$.*

- (i) *If $vs < v$, then $C^v \cap C_w \simeq C^{vs} \cap C_{ws}$.*
- (ii) *If $vs > v$ and $vs \not\leq ws$, then $C^v \cap C_w \simeq C^v \cap C_{ws} \times \mathbf{C}^*$.*
- (iii) *If $vs > v$ and $vs \leq ws$, then there exists a closed immersion*

$$(C^{vs} \cap C_{ws}) \times \mathbf{C} \hookrightarrow C^v \cap C_w$$

with open complement isomorphic to $(C^v \cap C_{ws}) \times \mathbf{C}^$.*

Corollary 2. *$\text{Hom}(\Delta_v, \Delta_w)$ is pure of weight 0.*

Proof. It is known that $\dim \operatorname{Hom}(\Delta_x, \Delta_y) = 1$ if and only if $x \leq y$, and that $\operatorname{Hom}(\Delta_x, \Delta_y) = 0$ if $x \not\leq y$. In particular, we may assume $v \leq w$ and proceed by induction on w . If $w = v$, the result is clear. Otherwise, pick a simple reflection s such that $ws < w$. In the situation of Proposition 1(i), the result is obvious. In the situation of Proposition 1(ii) and/or (iii), the long exact sequence of cohomology and Künneth formula yield an inclusion $\operatorname{Hom}(\Delta_v, \Delta_{ws}) \hookrightarrow \operatorname{Hom}(\Delta_v, \Delta_w)$. Since in both of these situations $v \leq ws$, this inclusion must be an isomorphism. \square

Corollary 3. *Let $v, w \in W$ with $v \leq w$. Let $s \in W$ be a simple reflection such that $ws > w$.*

- (i) *If $vs < v$, then $\operatorname{Ext}^1(\Delta_v, \Delta_{ws}) \simeq \operatorname{Ext}^1(\Delta_{vs}, \Delta_w)$.*
- (ii) *If $vs > v$ and $vs \not\leq w$, then $\operatorname{Ext}^1(\Delta_v, \Delta_w) \oplus \mathbf{Q}^H(-1) \simeq \operatorname{Ext}^1(\Delta_v, \Delta_{ws})$.*
- (iii) *If $vs > v$ and $vs \leq w$, then there is an exact sequence*

$$0 \rightarrow \mathbf{Q}^H(-1) \rightarrow \operatorname{Ext}^1(\Delta_v, \Delta_w) \oplus \mathbf{Q}^H(-1) \rightarrow \operatorname{Ext}^1(\Delta_v, \Delta_{ws}) \rightarrow \operatorname{Ext}^1(\Delta_{vs}, \Delta_w)(-1).$$

Proof. (i) is clear. Further, $\dim \operatorname{Hom}(\Delta_x, \Delta_y) = 1$ if and only if $x \leq y$. Thus, the Künneth formula yields (ii). In the situation of (iii):

$$\operatorname{Hom}(\Delta_v, \Delta_w) = \operatorname{Hom}(\Delta_v, \Delta_{ws}) = \operatorname{Hom}(\Delta_{vs}, \Delta_w) = \mathbf{Q}^H.$$

So the cohomology long exact sequence and Künneth formula yield (iii). \square

Theorem 4. *$\operatorname{Ext}^1(\Delta_v, \Delta_w)$ is pure of weight 2.*

Proof. We may assume $v \leq w$. Proceed by downwards induction on w . If w is the longest element, this is [M, Theorem 32]. If w is not the longest element, pick a simple reflection s such that $ws > w$ and apply Corollary 3. \square

Corollary 5. *Let $v, w \in W$ with $v \leq w$. Let $s \in W$ be a simple reflection such that $ws < w$. Then*

$$\dim \operatorname{Ext}^1(\Delta_v, \Delta_w) = \begin{cases} \dim \operatorname{Ext}^1(\Delta_{vs}, \Delta_{ws}) & \text{if } vs < v; \\ 1 + \dim \operatorname{Ext}^1(\Delta_v, \Delta_{ws}) & \text{if } vs > v \text{ and } vs \not\leq ws; \\ \dim \operatorname{Ext}^1(\Delta_v, \Delta_{ws}) & \text{if } vs > w \text{ and } vs \leq ws. \end{cases}$$

Some concluding observations:

- (i) $\dim \operatorname{Ext}^1(\Delta_v, \Delta_w)$ coincides with $(-1)^{\ell(w)-\ell(v)}$ times the coefficient of q in the corresponding Kazhdan-Lusztig R -polynomial.
- (ii) Theorem 4 implies that the graded algebra $\bigoplus_{v,w \in W} \operatorname{Ext}^\bullet(\Delta_v, \Delta_w)$ cannot, in general, be generated in degree 0 and 1. As otherwise $\operatorname{Ext}^i(\Delta_v, \Delta_w)$ would be pure of weight $2i$. This would contradict [Boe], since the Kazhdan-Lusztig R -polynomials are the Hodge-Euler polynomials of the $C^v \cap C_w$.

(iii) Using Corollary 5 one can check

$$\dim \operatorname{Ext}^1(\Delta_v, \Delta_w) = \dim \operatorname{Ext}^{\ell(w) - \ell(v) - 1}(\Delta_v, \Delta_w).$$

This upgrades to a canonical isomorphism by combining Theorem 4 with the main result of [BGS].

REFERENCES

- [BGS] A. BEILINSON, V. GINZBURG, W. SOERGEL, *Koszul duality patterns in Representation Theory*, Journal of A. M. S. **9** (1996), 473-527.
- [BGG] J. BERNSTEIN, I. M. GELFAND, S. I. GELFAND, *A certain category of \mathfrak{g} -modules*, Functional Analysis and its Applications **10**, no. 2 (1976), 1-8.
- [Boe] B. D. BOE, *A counterexample to the Gabber-Joseph conjecture*, Kazhdan-Lusztig theory and related topics (Chicago, IL, 1989), 1-3, Contemp. Math. **139**, Amer. Math. Soc., Providence, RI (1992).
- [M] V. MAZORCHUK, *Some homological properties of the category \mathcal{O}* , Pacific J. Math. **232** (2007), no.2, 313-341.
- [R] R. W. RICHARDSON, *Intersections of double cosets in algebraic groups*, Indag. Math. **3** (1992), 69-77.
- [RSW] S. RICHE, W. SOERGEL, G. WILLIAMSON, *Modular Koszul duality*, arXiv:1209.3760v1.

E-mail address: rsvirk@gmail.com